



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2007

Selenium-Containing Heterocycles From Isoselenocyanates: Synthesis of Ethyl 4-Oxo-2-amino-4,5-dihydroselenophene-3-carboxylates

Sommen, G L ; Linden, Anthony ; Heimgartner, H

Abstract: Aryl and alkyl isoselenocyanates 2 react with ethyl -chloroacetoacetate (1) in the presence of triethylamine to give the corresponding 4-oxo-2-amino-4,5-dihydroselenophene-3-carboxylates 4a-f in moderate to good yields.

DOI: <https://doi.org/10.2174/157017807780037342>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-50685>

Journal Article

Accepted Version

Originally published at:

Sommen, G L; Linden, Anthony; Heimgartner, H (2007). Selenium-Containing Heterocycles From Isoselenocyanates: Synthesis of Ethyl 4-Oxo-2-amino-4,5-dihydroselenophene-3-carboxylates. *Letters in Organic Chemistry*, 4(1):7-12.

DOI: <https://doi.org/10.2174/157017807780037342>

Selenium-Containing Heterocycles From Isoselenocyanates: Synthesis of Ethyl 4-Oxo-2-amino-4,5-dihydroselenophene-3-carboxylates

Geoffroy L. Sommen,¹⁾ Anthony Linden, Heinz Heimgartner*

Institute of Organic Chemistry, University of Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

Abstract: Aryl and alkyl isoselenocyanates **2** react with ethyl γ -chloroacetoacetate (**1**) in the presence of triethylamine to give the corresponding 4-oxo-2-amino-4,5-dihydroselenophene-3-carboxylates **4a-f** in moderate to good yields.

Key Words: Selenium heterocycles, Isoselenocyanates, Selenophenes, Cyclization, β -Ketoester, Crystal structure


INTRODUCTION

Selenium containing heterocycles are of increasing interest because of the antitumor, antibacterial and other biological activities of some representatives [1]. During our efforts devoted to the chemistry of selenaheterocycles, we also became interested in the synthesis of selenophenes. There are many articles and reviews dealing with the preparation of selenophenes [2], however, to the best of our knowledge, no synthesis starting from isoselenocyanates has been published. On the other hand, several procedures concerning the preparation of thiophenes by using isothiocyanates have been described [3]. For example, syntheses of some 4-oxothiophenes have been reported by Faull and Hull [4] as well as by Ibrahim, Sadek, Aziz, and Elnagdi [5] in the 80's. Compounds of this type, including 4-oxofurans, are known as novel antiallergic agents [6].

* Address correspondence to this author at the Institute of Organic Chemistry, University of Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland; E-mail: heimgart@oci.unizh.ch; ¹⁾ Present address: Lonza AG, Walliser Werke, CH-3930, Visp, Switzerland; e-mail: geoffroy.sommen@lonza.com

As a part of our program aimed at the development of simple procedures for the preparation of selenium-containing heterocycles [7-14], we have recently reported on the utility of isoselenocyanates for the synthesis of selenazetidines [15], selenazolidines [16,17], selenazines [16,18], selenadiazines [19], selenazepines [20], selenorhodanines [21], and triazoleselones [22]. As a continuation of this work, we report here on a novel and efficient synthesis of selenophene derivatives.

It was expected that the carbanion generated from ethyl γ -chloroacetoacetate (**1**) and triethylamine in DMF reacts with aromatic and aliphatic isoselenocyanates **2** to give intermediate keten-N,Se-acetals of type **3**. The latter, in an intramolecular nucleophilic substitution, can cyclize to give the selenophene ring system **4** (*Scheme 1*). Alternatively, ring closure via nucleophilic substitution by the N-atom could lead to pyrrolidine derivatives of type **5**. Analogous ring closure reactions, which gave azoleselones, have been described earlier [21,22]. The necessary isoselenocyanates **2** can be prepared easily from the corresponding *N*-arylformamides by treatment with phosgene and elemental selenium by following a slightly modified Barton procedure [23].









Scheme 1

RESULTS AND DISCUSSION

After stirring an equimolar mixture of ethyl γ -chloroacetoacetate (**1**), the respective isoselenocyanate **2**, and triethylamine in DMF at room temperature for four hours, the solvent was

evaporated. Column chromatography of the solid residue on silica gel with hexane/ethyl acetate and recrystallization of the major product from ethyl acetate gave ethyl 2-amino-4,5-dihydro-4-oxoselenophene-3-carboxylates **4** in fair yields (Table 1).

Table 1: Preparation of Selenophenes **4 from Isoselenocyanates **2****

Entry	Selenophenones 4	M.p. (°C)	Yield (%)
a		152–154	52
b		151–153	56
c		186–188	54
d		163–165	26
e		169–171	51
f		Oil	44

The structures of the products were established on the basis of their spectroscopic data and elemental analyses. Indicative are the IR absorptions at ca. 1660 and 1640 cm^{-1} for the keto and ester $\text{C}=\text{O}$ groups, respectively. In the mass spectra (CI mode), the respective $[\text{M}+1]^+$ peak appears as the base peak (100%), and the ^1H NMR spectra show the expected signals. Most informative are the ^{13}C NMR spectra with three signals for $\text{sp}^2\text{-C}$ atoms at ca. 193, 182, and 167 ppm for $\text{C}(4)=\text{O}$, $\text{C}(2)$, and CO_2Et , respectively. The atom $\text{C}(3)$ absorbs at ca. 99 ppm. In the case of **4b**, an X-ray crystal-structure determination was carried out, which confirmed the proposed structure (Fig. 1).




Fig. (1). ORTEP plot [24] of the molecular structure of **4b** (50% probability ellipsoids; arbitrary numbering of atoms)

In the crystal structure of **4b** all atoms of the molecule, except for the atoms of the 4-methylphenyl ring, form a planar system. When calculated using all atoms except $\text{C}(8)$ to $\text{C}(13)$, the maximum deviation from the plane is 0.054(2) Å for $\text{C}(17)$. The plane of the 4-methylphenyl ring lies perpendicular to the heterocyclic ring plane [angle: 88.94(7)°]. The amine group $\text{HN}(6)$ forms bifurcated hydrogen bonds. One interaction is intramolecular with the ester carbonyl $\text{O}(15)$ atom, while the second interaction is intermolecular with the same O -atom of a neighboring molecule, thereby linking pairs of molecules about a centre of inversion into dimeric moieties. The intramolecular interaction forms a closed loop with the graph set motif [25] of $\text{S}(6)$, while the intermolecular interaction

creates a loop with the graph set motif of $R^2_2(12)$. Taking both interactions together generates a four-membered loop with a binary graph set motif of $R^2_2(4)$.

In summary, we have evidenced that ethyl γ -chloroacetoacetate (**1**) is a very good precursor of selenophenes **4**. The base-catalyzed reaction with isoselenocyanates **2** proceeds *via* addition of the carbanion of **1** onto the isoselenocyanate to yield the acyclic intermediate **3**, which then undergoes a cyclization via the selenium atom to give **4** via elimination of HCl. The alternative ring closure via nucleophilic attack of the nitrogen atom of **3** (cf. [21,22]), which would lead to a pyrrole derivative, was not observed as a result of the higher nucleophilicity of the selenium atom. Thus, the selenophenes **4** have been easily prepared in a one-pot procedure starting from aryl and cyclohexyl isoselenocyanates. These building blocks are less toxic, relatively stable, safe and easy to handle, and can be synthesized conveniently.

EXPERIMENTAL

General

TLC: silica gel 60 F₂₅₄ plates (0.25 mm; Merck). Column chromatography (CC): silica gel 60 (0.040-0.063 mm; Merck). Melting Points: Büchi B-540 apparatus, in capillaries; uncorrected. ¹H-NMR (300 MHz) and ¹³C-NMR (75.5 MHz) spectra: Bruker ARX-300 instrument in CDCl₃; chemical shifts in ppm. CI-MS: Finnigan SSQ-700 or MAT-90 instrument; NH₃ as carrier gas.

Starting materials

Malononitrile and all halogenated compounds are commercially available (Fluka). Isoselenocyanates were prepared according to Barton's procedure [23] by starting from formamides. Formanilide and *N*-cyclohexylformamide are commercial products (Fluka and Aldrich), *N*-(4-methylphenyl)-, *N*-(4-bromophenyl)-, *N*-(2,6-dimethylphenyl)- and *N*-(4-methoxyphenyl)formamide were prepared from the respective aniline and 95% formic acid [26]. The solution was heated to reflux for 30 min and evaporated to dryness in vacuo. The residue was dissolved in ether and washed with diluted acetic acid (5%), water, and aqueous NaHCO₃ (5% aq.). The aqueous layer was extracted with ether, the combined organic extracts were dried with magnesium sulphate and evaporated under reduced pressure. The crude products were purified by recrystallization in EtOH/water.

General Procedure for the Preparation of 4,5-Dihydroselenophene-3-carboxylates 4a-f

A 25 mL round-bottom flask equipped with magnetic stirrer and condenser was charged with a solution of ethyl γ -chloroacetoacetate (**1**, 0.14 mL, 1.03 mmol) in DMF (20 mL). Triethylamine (0.14 mL, 1.03 mmol) was added and the mixture was stirred for 45 min at room temperature. Isoselenocyanate (**2**, 1.03 mmol) was added and the mixture was stirred for 4 h at room temperature before being evaporated to dryness under reduced pressure. The crude product was purified by column chromatography on silica gel without any further treatments by using hexane/ethyl acetate (100/0 to 50/50) as the eluant and crystallized from ethyl acetate.

Ethyl 4,5-dihydro-4-oxo-2-(phenylamino)selenophene-3-carboxylate (4a)

Yield: 52%. White crystals. M.p. 152-154°C. IR (KBr): 3441 w *br*, 3052 w , 2983 w , 1645 s , 1588 w , 1550 s , 1498 w , 1471 w , 1410 m , 1381 m , 1351 w , 1282 w , 1219 m , 1036 m , 946 w , 780 w , 736 w . ^1H NMR (CDCl_3): 1.65 (t, $J=7.1$ Hz, 3H, CH_3); 3.96 (s, 2H, CH_2); 4.61 (q, $J=7.1$ Hz, 2H, CH_2O); 7.62-7.73 (m, 5H, arom. CH). ^{13}C NMR (CDCl_3): 14.3 (CH_3); 32.6 (CH_2); 60.4 (CH_2O); 99.8 (C(3)); 124.1 (2 arom. CH); 128.0 (1 arom. CH); 129.7 (2 arom. CH); 138.1 (1 arom. C); 167.1 (CO_2); 181.9 (C(2)); 193.2 (CO). CI-MS: 314 (20), 313 (15), 312 (100, $[\text{M}+1]^+$), 311 (8, M^+), 310 (52), 309 (19), 308 (19). Anal. Calcd for $\text{C}_{13}\text{H}_{13}\text{NO}_3\text{Se}$ (311.01): C, 50.33; H, 4.22; N, 4.52. Found: C, 50.43; H, 4.38; N, 4.50.

Ethyl 4,5-dihydro-2-[(4-methylphenyl)amino]-4-oxoselenophene-3-carboxylate (4b)

Yield: 56%. White crystals. M.p. 151-153°C. IR (KBr): 3425 w *br*, 3224 w , 1662 s , 1636 s , 1565 s , 1513 m , 1470 w , 1416 m , 1374 m , 1224 s , 1190 w , 1043 m , 780 w . ^1H NMR (CDCl_3): 1.55 (t, $J=7.2$ Hz, 3H, CH_3); 2.55 (s, 3H, CH_3); 3.85 (s, 2H, CH_2); 4.52 (q, $J=7.1$ Hz, 2H, CH_2O); 7.40 (s, 4H, arom. CH). ^{13}C NMR (CDCl_3): 14.3 (CH_3); 21.0 (CH_3); 32.6 (CH_2); 60.4 (CH_2O); 99.6 (C(3)); 124.2 (2 arom. CH); 130.2 (2 arom. CH); 135.6 (1 arom. C); 138.2 (1 arom. C); 167.1 (CO_2); 182.2 (C(2)); 193.2 (CO). CI-MS: 328 (18), 327 (9), 326 (100, $[\text{M}+1]^+$), 325 (19, M^+), 324 (50), 323 (23), 322 (22). Anal. Calcd for $\text{C}_{14}\text{H}_{15}\text{NO}_3\text{Se}$ (325.02): C, 51.86; H, 4.66; N, 4.32. Found: C, 51.95; H, 4.83; N, 4.31.

Suitable crystals for the X-ray crystal-structure determination were grown from CH_2Cl_2 by slow evaporation of the solvent.

Ethyl 2-[(4-bromophenyl)amino]-4,5-dihydro-4-oxoselenophene-3-carboxylate (4c)

Yield: 54%. Yellowish crystals. M.p. 186-188°C. IR (KBr): 3442 w *br*, 3142 w , 1658 s , 1640 s , 1555 s , 1529 m , 1489 w , 1410 m , 1377 m , 1349 w , 1269 w , 1217 m , 1066 w , 1035 w , 1011 w , 782 w . ^1H NMR (CDCl_3): 1.33 (t, $J=7.2$ Hz, 3H, CH_3); 3.65 (s, 2H, CH_2); 4.30 (q, $J=7.1$ Hz, 2H, CH_2O); 7.19 (d, $J=8.2$

Hz, 2H, arom. CH); 7.52 (d, $J=8.2$ Hz, 2H, arom. CH). ^{13}C NMR (CDCl_3): 14.3 (CH_3); 32.8 (CH_2); 60.5 (CH_2O); 100.1 (C(3)); 121.6 (1 arom. C); 125.8 (2 arom. CH); 132.9 (2 arom. CH); 137.1 (1 arom. C); 167.2 (CO_2); 182.1 (C(2)); 193.0 (CO). CI-MS: 392 (78), 391 (34), 390 (100, $[\text{M}+1]^+$), 389 (23, M^+), 388 (47), 387 (19). Anal. Calcd for $\text{C}_{13}\text{H}_{12}\text{NO}_3\text{SeBr}$ (388.92): C, 40.13; H, 3.11; N, 3.60. Found: C, 40.31; H, 3.24; N, 3.56.

Ethyl 4,5-dihydro-2-[(2,6-dimethylphenyl)amino]-4-oxoselenophene-3-carboxylate (4d)

Yield: 26%. White crystals. M.p. 163-165°C. IR (KBr): 3441 w *br*, 3202 w , 2979 w , 1661 s , 1641 s , 1593 w , 1549 s , 1474 w , 1404 m , 1378 w , 1352 w , 1273 w , 1219 m , 1029 m , 790 w . ^1H NMR (CDCl_3): 1.33 (t, $J=7.2$ Hz, 3H, CH_3); 2.21 (s, 6H, 2 CH_3); 3.59 (s, 2H, CH_2); 4.28 (q, $J=7.1$ Hz, 2H, CH_2O); 7.07 (d, $J=8.2$ Hz, 2H, arom. CH); 7.20 (t, $J=8.2$ Hz, 1H, arom. CH); 10.98 (br s, 1H, NH). ^{13}C NMR (CDCl_3): 14.3 (CH_3); 17.8 (2 CH_3); 32.4 (CH_2); 60.2 (CH_2O); 99.0 (C(3)); 128.7 (2 arom. CH); 129.3 (arom. CH); 135.9 (1 arom. C); 136.1 (2 arom. C); 166.9 (CO_2); 185.2 (C(2)); 193.2 (CO). CI-MS: 342 (19), 341 (8), 340 (100, $[\text{M}+1]^+$), 339 (21, M^+), 338 (54), 337 (19), 336 (19). Anal. Calcd for $\text{C}_{13}\text{H}_{12}\text{NO}_3\text{SeBr}$ (339.04): C, 53.26; H, 5.07; N, 4.14. Found: C, 53.20; H, 5.27; N, 3.97.

Ethyl 4,5-dihydro-2-[(4-methoxyphenyl)amino]-4-oxoselenophene-3-carboxylate (4e)

Yield: 51%. White crystals. M.p. 169-171°C. IR (KBr): 3441 w *br*, 1638 s , 1607 w , 1544 s , 1511 m , 1461 w , 1385 w , 1336 w , 1240 w , 1212 m , 1183 w , 1028 m , 828 w , 786 w . ^1H NMR (CDCl_3): 1.45 (t, $J=7.1$ Hz, 3H, CH_3); 3.72 (s, 2H, CH_2); 3.89 (s, 3H, CH_3O); 4.41 (q, $J=7.1$ Hz, 2H, CH_2O); 7.01 (d, $J=8.2$ Hz, 2H, arom. CH); 7.32 (d, $J=8.2$ Hz, 2H, arom. CH); 11.54 (br s, 1H, NH). ^{13}C NMR (CDCl_3): 14.3 (CH_3); 32.6 (CH_2); 55.4 (CH_3O); 60.3 (CH_2O); 99.4 (C(3)); 114.7 (2 arom. CH); 126.2 (2 arom. CH); 130.9 (1 arom. C); 159.3 (1 arom. C); 167.1 (CO_2); 183.0 (C(2)); 193.2 (CO). CI-MS: 344 (8), 343 (33), 342 (100, $[\text{M}+1]^+$), 341 (12, M^+), 340 (42), 339 (10), 338 (9). Anal. Calcd for $\text{C}_{14}\text{H}_{15}\text{NO}_4\text{Se}$ (341.02): C, 49.42; H, 4.44; N, 4.12. Found: C, 49.66; H, 4.62; N, 4.02.

Ethyl 2-cyclohexylamino-4,5-dihydro-4-oxoselenophene-3-carboxylate (4f)

Yield: 44%. Orange oil. IR (film): 3173 w *br*, 2933 s , 2856 m , 1729 m *br*, 1642 s *br*, 1561 s , 1465 m , 1451 m , 1408 s , 1382 m , 1350 m , 1304 m , 1268 m , 1218 m , 1149 w , 1094 w , 1032 m , 916 w , 891 w , 858 w , 790 m , 732 m . ^1H NMR (CDCl_3): 1.19-1.47 (m, 7H, CH_3 , 2 CH_2); 1.95-1.99 (m, 2H, CH_2); 1.58-1.74 (m, 4H, 2 CH_2); 3.65 (s, 2H, CH_2); 3.26-3.31 (m, 1H, CH); 4.23 (q, $J=7.1$ Hz, 2H, CH_2O); 10.11 (br s, 1H, NH). ^{13}C NMR (CDCl_3): 14.3 (CH_3); 24.1 (2 CH_2); 24.9 (CH_2); 32.5 (2 CH_2); 57.5 (CH); 60.0

(CH₂O); 87.0 (C(3)); 157.4 (C(2)); 167.3 (CO₂); 196.5 (CO). CI-MS: 320 (11), 319 (34), 318 (100, [M+1]⁺), 317 (8, M⁺), 316 (44), 315 (9), 314 (12). Anal. Calcd for C₁₃H₁₉NO₃Se (317.05): C, 49.37; H, 6.06; N, 4.43. Found: C, 49.46; H, 5.92; N, 4.22.

Crystal-Structure Determination [27]

A crystal of **4b**, obtained from CH₂Cl₂, was mounted on a glass fibre and used for a low-temperature X-ray structure determination. All measurements were made on a *Nonius KappaCCD* area-detector diffractometer [28] using graphite-monochromated MoK α radiation ($\lambda = 0.71073$ Å) and an *Oxford Cryosystems Cryostream 700* cooler. Data reduction was performed with *HKL Denzo* and *Scalepack* [29]. The intensities were corrected for Lorentz and polarization effects, and an absorption correction based on the multi-scan method [30] was applied. Equivalent reflections were merged. Data collection and refinement parameters are given in [31]. A view of the molecule is shown in *Figure 1*. The structure was solved by direct methods using *SIR92* [32], which revealed the positions of all non-hydrogen atoms. The non-hydrogen atoms were refined anisotropically. The amine H-atom was placed in the position indicated by a difference electron density map and its position was allowed to refine together with an isotropic displacement parameter. All remaining H-atoms were placed in geometrically calculated positions and refined using a riding model where each H-atom was assigned a fixed isotropic displacement parameter with a value equal to 1.2U_{eq} of its parent C-atom (1.5U_{eq} for the methyl groups). Refinement of the structure was carried out on F^2 using full-matrix least-squares procedures, which minimized the function $\sum w(F_o^2 - F_c^2)^2$. A correction for secondary extinction was applied. Neutral atom scattering factors for non-hydrogen atoms were taken from [33a], and the scattering factors for H-atoms were taken from [34]. Anomalous dispersion effects were included in F_c [35]; the values for f' and f'' were those of [33b]. The values of the mass attenuation coefficients are those of [33c]. All calculations were performed using the *SHELXL97* [36] program.

ACKNOWLEDGMENTS

We thank the analytical units of our institute for spectra and analyses. Financial support of this work by the *Dr. Helmut Legerlotz Foundation* (postdoctoral stay of G.L.S. at the University of Zürich, 8.2004 – 8.2005) and *F. Hoffmann-La Roche AG*, Basel, is gratefully acknowledged.

REFERENCES

- [1] Koketsu, M.; Ishihara, H. *Curr. Org. Chem.*, **2003**, 7, 175.
- [2] (a) Sommen, G. L. *Mini Rev. Org. Chem.* **2005**, 2, 375; (b) Shatz, J. *Science of Synthesis: Product Class 11: Selenophenes*, **2002**, 9, 423; (c) Magdesevia, N. N. *Adv. Heterocycl. Chem.*, **1970**, 12, 1; (d) Hörnfeldt, A.-B. *Adv. Heterocycl. Chem.*, **1982**, 30, 127.
- [3] (a) Sommen, G. L.; Comel, A.; Kirsch, G. *Synlett*, **2001**, 11, 1731; (b) Sommen, G. L.; Comel, A.; Kirsch, G. *Tetrahedron Lett.*, **2002**, 1, 237; (c) Augustin, M.; Rudolf, W. D.; Schmidt, U. *Tetrahedron*, **1976**, 32, 3055; (d) Rudolf, W. D.; Schierhorn, A.; Augustin, M. *Tetrahedron*, **1979**, 35, 551.
- [4] (a) Faull, A. W.; Hull, R. *J. Chem. Soc., Perkin Trans. I*, **1981**, 1078; (b) Faull, A. W.; Hull, R. *J. Chem. Res.(S)*, **1979**, 240.
- [5] Ibrahim, N. S.; Sadek, K. U.; Aziz, S. I.; Elnagdi, M. H. *Z. Naturforsch.*, **1985**, 40B, 129.
- [6] Mack, R. A.; Zazulak, W. I.; Radov, L. A.; Baer, J. E.; Stewart, J. D.; Elzer, P. H.; Kinsolving, C. R.; Georgiev, V. S. *J. Med. Chem.*, **1988**, 31, 1910.
- [7] Atanassov, P. K.; Linden, A.; Heimgartner, H. *Helv. Chim. Acta*, **2004**, 87, 1873.
- [8] Atanassov, P. K.; Linden, A.; Heimgartner, H. *Helv. Chim. Acta*, **2004**, 87, 1452.
- [9] Atanassov, P. K.; Linden, A.; Heimgartner, H. *Heterocycles*, **2004**, 62, 521.
- [10] Atanassov, P. K.; Linden, A.; Heimgartner, H. *Heterocycles*, **2003**, 61, 569.
- [11] Atanassov, P. K.; Linden, A.; Heimgartner, H. *Helv. Chim. Acta*, **2003**, 86, 3235.
- [12] Atanassov, P. K.; Zhou, Y.; Linden, A.; Heimgartner, H. *Helv. Chim. Acta* **2002**, 85, 1102.
- [13] Zhou, Y.; Heimgartner, H. *Helv. Chim. Acta*, **2000**, 83, 539.
- [14] Zhou, Y.; Linden, A.; Heimgartner, H. *Helv. Chim. Acta*, **2000**, 83, 1576.
- [15] Sommen, G. L.; Linden, A.; Heimgartner, H. *Helv. Chim. Acta*, **2005**, 88, 766.
- [16] Sommen, G. L.; Linden, A.; Heimgartner, H. *Eur. J. Org. Chem.*, **2005**, 3128.
- [17] Sommen, G. L.; Linden, A.; Heimgartner, H. *Tetrahedron*, **2006**, 62, 3344.
- [18] Sommen, G. L.; Linden, A.; Heimgartner, H. *Helv. Chim. Acta*, accepted.
- [19] Sommen, G. L.; Linden, A.; Heimgartner, H. *Helv. Chim. Acta*, **2006**, 89, 1322.
- [20] Sommen, G. L.; Linden, A.; Heimgartner, H. *Tetrahedron Lett.*, **2005**, 46, 6723.
- [21] Sommen, G. L.; Linden, A.; Heimgartner, H. *Heterocycles*, **2005**, 65, 1903.
- [22] Favero, F.; Sommen, G. L.; Linden, A.; Heimgartner, H. *Heterocycles*, **2006**, 67, 749.

- [23] (a) Barton, D. H. R.; Parekh, S. I.; Tajbakhsh, M.; Theodorakis, E. A.; Tse, C.-L. *Tetrahedron*, **1994**, *50*, 639; (b) Bakhsh, M. T.; Behshtiha, Y. S.; Heravi, M. M. *J. Chem. Soc. Pakistan*, **1996**, *18*, 159.
- [24] Johnson, C. K. *ORTEP II*, Report ORNL-5138, Oak Ridge National Laboratory, Oak Ridge, Tennessee, 1976.
- [25] Bernstein, J.; Davis, R. E.; Shimon, L.; Chang, N.-L. *Angew. Chem. Int. Ed. Engl.*, **1995**, *34*, 1555.
- [26] Katritzky, A. R.; Parris, R. L.; Ignatchenko, E. S.; Allin, S. M.; Siskin, M. *J. Prakt. Chem.* **1997**, *339*, 59.
- [27] CCDC-629496 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre CCDC via http://www.ccdc.ac.uk/data_request/cif.
- [28] Hooft, R. *KappaCCD Collect Software*, Nonius BV, Delft, The Netherlands, **1999**.
- [29] Otwinowski, Z., Minor, W. in *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, Eds. Carter, C. W. Jr.; Sweet, R. M., Academic Press, New York, **1997**, 307.
- [30] Blessing, R.H. *Acta Crystallogr., Sect A* **1995**, *51*, 33.
- [31] Crystallographic data for compound **4b**: Empirical formula: C₁₄H₁₅NO₃Se; formula weight: 324.18 g mol⁻¹; crystal color, habit: colorless, prism; crystal dimensions: 0.20 × 0.30 × 0.30 mm; temperature: 160(1) K; crystal system: monoclinic; space group: *C2/c*; *Z* = 8; reflections for cell determination: 28982; 2 θ range for cell determination: 4 – 60°; unit cell parameters: *a* = 26.8226(4) Å, *b* = 7.2506(1) Å, *c* = 14.1780(2) Å, β = 97.811(1)°, *V* = 2731.76(7) Å³, *D_x* = 1.576 g cm⁻³; $\mu(\text{MoK}\alpha)$ = 2.750 mm⁻¹; scan type: ϕ and ω ; 2 $\theta_{(\text{max})}$ = 60°; transmission factors (min; max): 0.483; 0.581; total reflections measured: 41421; symmetry independent reflections: 3990; reflections with *I* > 2 σ (*I*): 3431; reflections used in refinement: 3990; parameters refined: 179; final *R*(*F*) [*I* > 2 σ (*I*) reflections]: 0.0319; *wR*(*F*²) (all data): 0.0820; weights: $w = [\sigma^2(F_o^2) + (0.0331P)^2 + 3.6031P]^{-1}$ where $P = (F_o^2 + 2F_c^2) / 3$; goodness of fit: 1.136; secondary extinction coefficient: 0.0012(2); final $\Delta_{\text{max}}/\sigma = 0.001$; $\Delta\rho$ (max; min) = 0.46; -0.60 [e Å⁻³].
- [32] Altomare, A.; Cascarano, G.; Giacovazzo, C.; Guagliardi, A.; Burla, M. C.; Polidori, G.; Camalli, M. *SIR92, J. Appl. Crystallogr.* **1994**, *27*, 435.

- [33] (a) Maslen, E. N.; Fox, A. G.; O'Keefe, M. A. in 'International Tables for Crystallography', Ed. Wilson, A. J. C., Kluwer Academic Publishers, Dordrecht, **1992**, Vol. C, Table 6.1.1.1, p. 477; b) Creagh, D. C.; McAuley, W. J. *ibid.* Table 4.2.6.8, p. 219; c) Creagh, D. C.; Hubbell, J. H. *ibid.* 4] Stewart, R. F.; Davidson, E. R.; Simpson, W. T. *J. Chem. Phys.* **1965**, 42, 3175.
- [35] Ibers, J. A.; Hamilton, W. C. *Acta Crystallogr.* **1964**, 17, 781.
- [36] Sheldrick, G. M. *SHELXL97, Program for the Refinement of Crystal Structures*, University of Göttingen, Germany, 1997.